

## INTERACTION OF *DZRI*, *OPAQUE-2* AND NORMAL ENDOSPERM MAIZE INBRED LINES FOR GRAIN YIELD AND PROTEIN QUALITY

Sekip ERDAL<sup>1\*</sup>, Ahmet OZTURK<sup>1</sup>, Rahime CENGIZ<sup>2</sup>, Mehmet PAMUKCU<sup>1</sup>, Cuneyt DINCER<sup>3</sup>, Bulent CENGIZ<sup>4</sup>

<sup>1</sup>Western Mediterranean Agricultural Research Institute, Antalya, TURKEY

<sup>2</sup>Sakarya University of Applied Sciences, Faculty of Agriculture, Department of Field Crops, Sakarya, TURKEY

<sup>3</sup>Akdeniz University, Finike Vocational School, Antalya, TURKEY

<sup>4</sup>Maize Research Institute, Sakarya, TURKEY

\*Corresponding author: [sekip65@yahoo.com](mailto:sekip65@yahoo.com)

Received: 19.08.2020

### ABSTRACT

The objectives of this study were to (i) investigate genetic relationships among high lysine (HK) (*opaque-2*), high methionine (HM) (*dzr1*) and high yielding maize inbred lines (ii) to evaluate grain yield and protein quality of hybrids produced from these germplasm groups. Fifty-six hybrids generated from an 8 × 8 full diallel mating design were tested at two locations in Turkey in 2017 and 2018. Significant reciprocal effects for lysine, methionine, lysine quality index and methionine quality index revealed that parent effects may not be ignored in breeding for these traits. Lower grain yield among crosses produced from non-normal endosperm suggested that for high yield and improved amino acid concentration at least one parent of HK or HM should be used in hybrid combinations. HM × HM hybrids were not only good for high methionine, but also high lysine and protein. Several of the experimental hybrids in this study outperformed the commercial checks in terms of lysine and methionine yield. M2 × S1 hybrid had 27 % more lysine (58.6 kg ha<sup>-1</sup>) and 26 % more methionine (42.5 kg ha<sup>-1</sup>) yield than commercial high yielding normal hybrids. This study revealed that HK and HM germplasm can be combined with adapted high yield maize inbred lines to develop high yielding, high methionine and high lysine hybrids.

**Keywords:** Combining ability, diallel analysis, hybrid, lysine, methionine

### INTRODUCTION

Maize protein is inadequate in some essential amino acids. Lysine and methionine are not sufficient to meet the nutritional requirements of monogastric animals, including humans (Schutte and Jong, 1999; Ravindran, 2012). Lysine and methionine are the two most important amino acids in maize-soybean poultry feed rations because costly supplementation is required to meet the recommended levels (Scott et al., 2008). Studies show that increasing methionine levels significantly increased egg production and egg weight of laying hens (Harms et al., 1998; Saki et al., 2012). In feeding trials, chickens and pigs fed high lysine maize had a higher body weight than those fed normal maize hybrids (Krivanek et al., 2007).

High lysine maize called QPM (Quality Protein Maize) contains high levels of lysine and tryptophan, the protein level is similar to normal maize. In these genotypes, amino acid balance is under the control of a recessive gene called *opaque-2* (*o2*) which is a natural

mutation (Mertz et al., 1964). Modifier/enhancer genes of the *o2o2o2* endosperm confer higher lysine in a phenotypically desirable kernel type (Vasal et al., 1993; Krivanek et al., 2007).

In a study by Phillips and McClure (1985), several lines with elevated methionine levels were identified including a line from the Iowa Stiff-Stalk Synthetic population designated BSSS53. The 10-kDa delta zein (*dzs10*) is elevated in this line, causing the high methionine concentration reported by Kirihara et al. (1988). Later it was understood that *dzs10* transcripts are regulated by a gene called delta zein regulator1 (*dzr1*). Olsen et al. (2003) successfully released high methionine versions of the public inbred lines A632, B73 and Mo17 containing *dzr1*.

A study carried out by Scott et al. (2004) revealed that methionine is reduced in *o2o2* maize germplasm. Compositional analysis of a set of QPM hybrids showed that QPM hybrids had lower methionine levels than even

normal maize varieties (Mbuya et al., 2011). These observations suggest that use of the *opaque-2* gene in maize in poultry feed might increase the methionine supplementation requirement. Huffman et al. (2016) used a diallel study to investigate interaction of genetic mechanisms between *floury-2* and *dzr1*. Interaction among *opaque-2* which is the most widely used lysine modification mutation, and *dzr1* may help in breeding for these traits. The objectives of this study are (i) to investigate genetic relationships among *dzr1* (high methionine), *opaque-2* (high lysine, QPM) and high yielding maize inbred lines based on Griffing's diallel analysis (ii) to determine grain yield and amino acid levels of the hybrids obtained from an 8 × 8 full diallel mating design using these three germplasm groups.

## MATERIALS AND METHODS

### Germplasm

The maize inbred lines used in the study are listed in Table 1. The lines K1 and K2 are rich in lysine (QPM) due to the *opaque-2* mutation (Betran et al., 2003a; Betran et al., 2003b). The lines M1, M2 and M3 are high methionine versions of A632, B73 and Mo17, respectively, and they reportedly contain 12.5%, 25% and 50% more methionine than their wild-type counterparts (Philips et al., 2008). The line A1 is an elite line developed by Bati Akdeniz Agricultural Research Institute with high yield and drought tolerance (Erdal, 2019). The lines S1 and S2 are high yielding commercial maize lines released by the Maize Research Institute of Turkey. Two widely grown high yielding normal endosperm commercial hybrids, P31G98 and DKC6589, were included in the study as checks. Crosses were done to make a complete (including reciprocals) eight by eight diallel mating design in 2016.

**Table 1.** Maize inbred lines used in the eight by eight diallel design

Line	Name	Specific trait	Heterotic group	Source
K1	T×802	High lysine ( <i>opaque-2</i> , QPM)	Tropical / subtropical	Texas A&M University
K2	T×807	High lysine ( <i>opaque-2</i> , QPM)	Tropical / subtropical	Texas A&M University
M1	58609 A632 (Meth)	High methionine ( <i>dzr1</i> )	Stiff-Stalk	Minnesota Agricultural Experiment Station
M2	58615 B73 (Meth)	High methionine ( <i>dzr1</i> )	Stiff-Stalk	Minnesota Agricultural Experiment Station
M3	58803 Mo17 (Meth)	High methionine ( <i>dzr1</i> )	Lancaster	Minnesota Agricultural Experiment Station
A1	Ant-24702	High yielding/elite/ drought tolerant	Lancaster	Bati Akdeniz Agricultural Research Institute
S1	ADK-451	High yielding /Commercial line	Leaming	Maize Research Institute of Turkey
S2	ADK-455	High yielding /Commercial line	Stiff-Stalk	Maize Research Institute of Turkey

### Experimental design

All 56 hybrids and the two check varieties were evaluated as entries in a randomized complete block design with three replications. The locations Antalya (southern Turkey) and Sakarya (northern Turkey) were used in 2017 and 2018 to produce four testing environments. Antalya (36°52'N 30° 45'E) is a typical mediterranean province where the climate is warm in summer and the winter months are more rainier than the summer months. Sakarya (40°48'N 30° 25'E) province has a rainy and humid weather and a temperate climate. Winters are rainy and warm, and summers are hot. Experimental plots consisted of 2 rows, 5 m long and 0.70 m between rows. In order to eliminate the pollen effect on kernel protein quality, 5 plants in the first row of each plot were self pollinated and the resulting grain was bulked for amino acid analysis. Sowing were done in April (Antalya

and May (Sakarya) in both years. Before sowing, 600 kg ha<sup>-1</sup> of composite 15-15-15 fertilizer was applied to the experiments area to provide 90 kg ha<sup>-1</sup> for each N, P and K elements. Later, 152 kg ha<sup>-1</sup> of N dose using 46% urea fertilizer, was given to the soil in a few different times (TTSM, 2018). To control weeds in the experiments, 225 g / L Isoxaflutole + 90 g / L Thiencazone-methyl + 150 g / L Cyprosulfamide active ingredient herbicide was applied after plant emergence. Experiments were regularly irrigated to avoid water stress till harvest.

Grain yield (t ha<sup>-1</sup>) was determined from the second row of each plot and adjusted to 15% grain moisture content at harvest. Protein concentration was determined by the Dumas classical method (AOAC International, 2002). Liquid Chromatography Triple Quadrupole Mass Spectrometry (LC-MS/MS) was used for quantifying methionine and lysine. Before analysis, maize samples

were hydrolyzed using the method of Chan and Matanjun (2017) which was modified according to Faountoukakakis and Lahm (1998). Amount of 0.2 g of the sample was homogenized and weighed into a solution of 10 mL of 6 N HCl (containing 0.02% of phenol). The mixture was mixed by vortexing in a tightly sealed test tube for 5 min and then stored in an oven at 110 °C for 24 h to complete the hydrolysis. Following cooling to room temperature, the mixture was filtered through a 0.45 µm PTFE membrane and then injected into the LC-MS/MS device. Total lysine and methionine values were calculated as concentration (g / 100 g, dry matter). Protein quality indexes were calculated for both lysine and methionine by dividing the amino acid concentration by the total protein concentration.

### Statistical analysis

The field plot data was subjected to analysis of variance (ANOVA). Least significant difference (LSD) test (Steel and Torrie, 1980) was used when the difference

between means were statistically significant. Analysis of Genetic Designs, AGD-R, a statistical software programme which was developed by Rodriguez et al. (2015) was used to analyze the diallel data using Griffing (1956) method III model I (fixed). In this analysis, variance was partitioned into the components listed in Table 2, which were all considered to be fixed effects in a linear model. The model effects for general combining ability (GCA), specific combining ability (SCA), reciprocal, maternal, non-maternal and their interactions were tested for significance using an F-test based on the estimated sums of squares generated by the linear model. In addition, variance components such as GCA / SCA ratio, phenotypic variance, narrow and broad sense heritability were estimated using AGD-R (Rodriguez et al., 2015). Comparisons between the high lysine (HK), high methionine (HM) and high yield (HY) groups were made using specific contrasts and trait correlations were determined using Pearson correlation coefficients.

**Table 2.** Analysis of variance (ANOVA) and variance components for investigated traits

Source of variation	DF	Grain yield (t ha <sup>-1</sup> )	Protein (%)	Lysine (g/100g)	Methionine (g/100g)	KQI (%)	MQI (%)
Mean square error							
Environment (E)	3	11082806 **	13.80 **	0.022182 **	0.00976 **	4.99 **	5.99 **
Replication	8	80162.84 **	1.53 **	0.000444 *	0.001928 **	0.04 *	0.84 **
Genotype (G)	55	257512.8 **	6.05 **	0.008789 **	0.004297 **	0.66 **	17.69 **
GCA	7	1492202 **	37.93 **	0.027756 **	0.018349 **	0.80 **	6.44 *
SCA	20	99774.21 *	1.88 *	0.005936 **	0.001595*	0.43 ns	3.33 ns
Reciprocal (R)	28	61511 ns	1.06 ns	0.006085 **	0.002715 **	0.79 **	7.92 **
Maternal (M)	7	103063.3 *	1.16 ns	0.014525 **	0.006551 *	1.76 **	3.36 ns
Non-maternal (NM)	21	47660.22 ns	1.03 ns	0.003272 **	0.001436 **	0.46 **	4.56 **
E×G	165	60459.03 **	1.24 **	0.001425 **	0.001213 **	0.19 **	23.17 **
E×GCA	21	119838.6 **	2.92 **	0.000701 ns	0.003915 **	0.08 ns	7.45 **
E×SCA	60	47310.73 **	1.07 **	0.001926 **	0.000881 **	0.26 **	7.09 **
E×R	84	55005.8 **	0.94 **	0.001248 **	0.000775 **	0.17 **	8.62 **
E×M	21	47423.91 ns	0.94 ns	0.002534 **	0.002004 **	0.25 *	4.74 **
E×NM	63	57533.09 **	0.94 **	0.00082 **	0.000366 **	0.14 **	3.88 **
Residual	440	30509.21	0.29	0.000156	0.000147	0.01	3.17
Magnitude of the variance components							
GCA		40602.57	1.05	0.00115	0.000758	0.03	0.048
SCA		11544.17	0.27	0.001445	0.000362	0.10	0.038
M		1154.23	0.01	0.000352	0.00016	0.04	0.008
NM		2858.50	0.12	0.000779	0.000322	0.11	0.051
GCA/SCA		3.52	3.94	0.80	2.09	0.32	0.99
Phenotypic Variance		128425.50	2.77	0.005384	0.002668	0.38	0.195
Narrow Heritability		0.63	0.75	0.43	0.57	0.17	0.39
Broad Heritability		0.72	0.85	0.70	0.70	0.45	0.58

\* Significant at the 0.05 probability level, \*\* Significant at the 0.01 probability level, ns not significant

## RESULTS AND DISCUSSION

### Analysis of variance and genetic components

Results of combined analysis of variance (ANOVA) and genetic components for measured traits are presented in Table 2. Since the differences among genotypes were significant, diallel analysis was performed. General combining ability (GCA) effects are considered to be governed by additive gene action and were found to be significant for all traits. Significant specific combining ability (SCA) effects were detected in grain yield,

protein, lysine, and methionine while non-significant effects were determined in lysine quality index (KQI) and methionine quality index (MQI). Significant reciprocal (R) effects were identified for lysine, methionine, KQI and MQI traits, showing that the direction of the cross impacted the cross performance. Therefore it would be beneficial to evaluate crosses and their reciprocals in test experiments. Significant maternal effects were found for grain yield, lysine, methionine and KQI. Significant environment (E) × GCA, E × SCA and E × R interactions

showed the importance of environmental effects on these genetic components (Table 2).

GCA/SCA ratios may give some information about the trait heritability, because the additive effects determine GCA are more heritable than the non-additive effects that determine SCA. Non-additive gene effects are more prevalent in lysine, KQI and MQI traits (GCA/SCA <1). The highest narrow sense heritability value (0.75) was for protein while the lowest value was obtained for KQI (0.17). Both narrow and broad sense heritability were lower for lysine and methionine when compared to grain yield and protein (Table 2).

Although grain yield is reportedly influenced by non-additive genetic variance (Nas et al., 2000) in maize, Duraes et al. (2002) Hallauer and Carena (2009) and Erdal et al. (2015) reported that the variance of additive gene effects was more important for grain yield. Heritability varies greatly among studies and is dependent on the genotypes used in a study, the environmental conditions and many other factors. In this study, relatively high heritability is expected due to presence of large effect genes controlling the traits of interest. For example, *o2* has a large effect on both lysine and yield. High narrow (0.75) and broad (0.85) sense heritability values suggest protein concentration can be improved via breeding. High additive gene effects for methionine revealed by Huffman et al. (2016) are consistent with our results related to methionine heritability.

#### Combining ability analysis

General combining ability (GCA) effects for investigated traits are presented in Table 3. Significant and positive GCA values for yield were obtained from high

yielding normal endosperm (S1 and A1) inbred lines. M1 (*dzr1*), K1 (*opaque-2*) and K2 (*opaque-2*) inbred lines have negative and significant effects indicating the lower yield potential of high amino acid germplasm. Grain protein concentration GCA values were statistically significant in all lines ( $p < 0.01$  and  $p < 0.05$ ). The highest and lowest results were obtained from the M1 (0.82) and S1 (-0.65) lines, respectively. The highest positive and significant lysine GCA effect was obtained from line M1 (0.016), while the lines K2, M3, M2 followed. Negative and significant GCA for lysine was obtained in normal endosperm lines (A1, S1, S2). When methionine GCA values were examined, it was determined that the lines containing *dzr1* (M1, M2 and M3) have the highest positive and significant results ( $p < 0.01$ ). Therefore, we conclude that *dzr1* lines are effective parents for creating high methionine hybrids. When lysine quality index (KQI) results were examined, positive and significant results were obtained only for K1 and K2 lines (QPM). It was determined that the S1, S2 (normal endosperm) and M2 lines were the best lines for MQI GCA effects.

Specific combining ability (SCA) effects for investigated traits are given in Table 4. The highest positive and significant SCA effects in terms of grain yield were from the combinations M1  $\times$  K1 (129.83), A1  $\times$  M3 (66.47) and M3  $\times$  K2 (64.07). In the grain protein concentration, A1  $\times$  M3 (0.46) ranked first while M1  $\times$  K1 combination (0.40) was second and S2  $\times$  K1 (0.36) was the third. The highest positive and significant SCA effects in terms of grain lysine concentration were from the M1  $\times$  K1, A1  $\times$  M2 and S2  $\times$  K1 combinations, respectively. S1  $\times$  M3, S1  $\times$  K2 and M1  $\times$  K1 were the best combinations for methionine. A1  $\times$  M2 for KQI and S1  $\times$  M3 were the most successful combinations in terms of MQI.

**Table 3.** General combining ability effects for grain yield, protein, lysine, methionine, lysine quality index (KQI) and methionine quality index (MQI)

Parents	Grain Yield	Protein	Lysine	Methionine	KQI	MQI
K1	-28.80*	-0.42**	-0.00071 ns	-0.01615**	0.151511**	-0.04975**
K2	-34.22*	0.11*	0.014494**	-0.01334**	0.078722**	-0.16689**
M1	-191.15**	0.82**	0.015601**	0.015951**	-0.14622**	-0.05706**
M2	-10.78ns	0.51**	0.013629**	0.021875**	-0.05899**	0.073152**
M3	1.23ns	0.27**	0.013807**	0.006157**	0.045614**	-0.0011ns
A1	98.68**	-0.18**	-0.01036**	-0.00867**	-0.0261*	-0.02949*
S1	155.06**	-0.65**	-0.02509**	-0.00716**	-0.00673 ns	0.099292**
S2	9.98ns	-0.46**	-0.02138**	0.001339 ns	-0.03782**	0.131847**

\* Significant at the 0.05 probability level, \*\* Significant at the 0.01 probability level, ns not significant

It was observed that M1, M2, M3 and K2 inbred lines can be used for high protein concentration breeding studies. Methionine germplasm did not only have good GCA effects for methionine but also for high lysine concentration. Higher KQI GCA effects of *opaque-2* lines are consistent with previous studies that have shown QPM hybrids have high protein quality in terms of lysine. The International Maize and Wheat Improvement Center (CIMMYT) is one of the leading institutions to develop QPM (high tryptophan and lysine) lines and hybrids (Vivek et al., 2008). CIMMYT breeders generally used

the KQI trait when developing QPM varieties. Since QPM lines used in our study were developed from CIMMYT populations, it was not surprising that these lines had high GCA values in terms of KQI. When the methionine quality index (MQI) GCA values were examined, it was observed that the first two ranks were from the S1 and S2 lines, which were high yielding and normal endosperm lines. These lines did not produce very good results in terms of methionine concentration, but they combined very well with any high methionine (M1, M2 and M3) line, in other words, they served as complementary lines.

Therefore, these two lines can be used for their complementary effects.

**Table 4.** Specific combining ability effects for grain yield, protein, lysine, methionine, lysine quality index (KQI) and methionine quality index (MQI)

Hybrid	Grain yield	Protein	Lysine	Methionine	KQI	MQI
K2 × K1	-93.34**	-0.09 <sup>ns</sup>	0.013146**	-0.0017 <sup>ns</sup>	0.157093**	-0.01073 <sup>ns</sup>
M1 × K1	129.83**	0.40**	0.031946**	0.010703**	0.152002**	0.014326 <sup>ns</sup>
M1 × K2	8.28 <sup>ns</sup>	-0.27**	0.005379*	0.000644 <sup>ns</sup>	0.112161**	0.06665*
M2 × K1	52.19 <sup>ns</sup>	-0.01 <sup>ns</sup>	-0.0324**	-0.0162**	-0.28243**	-0.12609**
M2 × K2	-27.59 <sup>ns</sup>	0.26**	-0.02323**	-0.00736**	-0.29872**	-0.12222**
M2 × M1	-114.95**	-0.01 <sup>ns</sup>	-0.00248 <sup>ns</sup>	0.000882 <sup>ns</sup>	0.007736 <sup>ns</sup>	0.015117 <sup>ns</sup>
M3 × K1	-67.68*	-0.51**	-0.01063**	0.002476 <sup>ns</sup>	0.07488**	0.135216**
M3 × K2	64.07*	0.16 <sup>ns</sup>	-0.00863**	-0.01608**	-0.13679**	-0.17493**
M3 × M1	35.29 <sup>ns</sup>	-0.20*	-0.0159**	-0.00847**	-0.08127**	-0.03794 <sup>ns</sup>
M3 × M2	-18.36 <sup>ns</sup>	0.12 <sup>ns</sup>	0.015736**	0.001421 <sup>ns</sup>	0.076402**	-0.03647 <sup>ns</sup>
A1 × K1	-37.31 <sup>ns</sup>	-0.28**	-0.02839**	-0.00246 <sup>ns</sup>	-0.17473**	0.046471 <sup>ns</sup>
A1 × K2	9.29 <sup>ns</sup>	-0.06 <sup>ns</sup>	-0.00663*	0.009444**	-0.04583 <sup>ns</sup>	0.098266**
A1 × M1	-78.34*	0.31**	0.007127**	0.006346*	-0.03874 <sup>ns</sup>	-0.01812 <sup>ns</sup>
A1 × M2	43.98 <sup>ns</sup>	-0.08 <sup>ns</sup>	0.031017**	0.006194*	0.303533**	0.063593*
A1 × M3	66.47*	0.46**	0.018261**	0.003095 <sup>ns</sup>	0.027155 <sup>ns</sup>	-0.06987**
S1 × K1	38.21 <sup>ns</sup>	0.13 <sup>ns</sup>	0.003883 <sup>ns</sup>	-0.0019 <sup>ns</sup>	-0.04924 <sup>ns</sup>	-0.08504**
S1 × K2	26.83 <sup>ns</sup>	0.21*	0.011974**	0.011587**	0.102396**	0.088927**
S1 × M1	-42.09 <sup>ns</sup>	-0.28**	-0.00564*	-0.01064**	0.046177 <sup>ns</sup>	-0.03273 <sup>ns</sup>
S1 × M2	62.04*	-0.13 <sup>ns</sup>	0.013501**	0.009233**	0.161581**	0.117808**
S1 × M3	-51.93 <sup>ns</sup>	-0.10 <sup>ns</sup>	-0.00621*	-0.013102**	-0.01968 <sup>ns</sup>	0.154996**
S1 × A1	-4.91 <sup>ns</sup>	-0.01 <sup>ns</sup>	-0.01183**	-0.01033**	-0.09369**	-0.08724**
S2 × K1	-21.91 <sup>ns</sup>	0.36**	0.02245**	0.009082**	0.122438**	0.025842 <sup>ns</sup>
S2 × K2	12.45 <sup>ns</sup>	-0.20*	0.007984**	0.003458 <sup>ns</sup>	0.109682**	0.054029*
S2 × M1	61.97*	0.05 <sup>ns</sup>	-0.02043**	0.000529 <sup>ns</sup>	-0.19807**	-0.00731 <sup>ns</sup>
S2 × M2	2.68 <sup>ns</sup>	-0.15 <sup>ns</sup>	-0.00215 <sup>ns</sup>	0.00583*	0.0319 <sup>ns</sup>	0.088261**
S2 × M3	-27.86 <sup>ns</sup>	0.08 <sup>ns</sup>	0.007378**	0.00445 <sup>ns</sup>	0.059298*	0.028989 <sup>ns</sup>
S2 × A1	0.82 <sup>ns</sup>	-0.33**	-0.00955**	-0.01229**	0.02295 <sup>ns</sup>	-0.03309 <sup>ns</sup>
S2 × S1	-28.15 <sup>ns</sup>	0.19*	-0.00568*	-0.01106**	-0.14754**	-0.15672**

\* Significant at the 0.05 probability level, \*\* Significant at the 0.01 probability level, <sup>ns</sup> not significant

#### Contrasts between germplasm groups

In order to better understand the effects of the major genes *dzr1* and *opaque-2* in the germplasm used in this study, we compared the performance of the three germplasm groups: HY (no major genes), HM (*dzr1*) and HK (*opaque-2*) (Table 5). The high yielding lines (HY) gave the highest grain yield values in both the male and female group comparisons. On the other hand, the high lysine (HK) and high methionine (HM) groups were found to have similar grain yields. HM lines were found to have the highest protein concentration in both female and male, while HY lines were found to have low protein values (Table 5).

When the interactions were considered, it was observed that combinations of HM × HK (0.367 g / 100 g) and HM × HM (0.362 g / 100g) gave the highest lysine results, while the lowest result was produced in the HY × HY combination (0.287 g / 100g) (Table 5).

The highest methionine concentrations were obtained from the HM group and the highest lysine concentrations were obtained from the HK group regardless of which parent contributed the germplasm group. The HK group had lowest methionine levels regardless of which parent contributed this group. Interestingly, the HM × HM combination was in the top significance group for both

lysine and methionine concentrations, while the HK × HK combination was in the top significance group for lysine concentration, but in the bottom significance group for methionine.

The contrast between the three germplasm groups HY, HK and HM showed that lowest yield results were obtained from hybrids of HK × HK and HM × HM lines. This finding shows that quality protein (both methionine and lysine) germplasm that we used in this study have low yield potential. Using HM and HK germplasm classes in combination with HY groups could be an option. However, this is rarely done in protein quality breeding programs. It is therefore of interest to consider our results in light of what is known about the gene action of these major genes. Although lysine levels are influenced by the *opaque-2* gene, the level of lysine in genotypes varies due to modifier / enhancer loci in QPM genetic material. This supports the suggestion that lysine levels should be measured throughout the course of breeding programs (Vivek et al., 2008). Our data suggests that *dzr1* may be a modifier of lysine levels when used as a female parent. While HK × HK hybrids had high lysine levels, they had low methionine levels, consistent with previous reports that *opaque-2* hybrids have low methionine levels Scott et al. (2004).

**Table 5.** Effect of parents on grain yield (t ha<sup>-1</sup>), protein (%), lysine (g/100g) and methionine (%) concentration

	Female group			
	Yield**	Protein**	Lysine**	Methionine**
HK	11.69 b	10.26 b	0.336 b	0.212 c
HM	11.99 b	10.91 a	0.359 a	0.253 a
HY	13.35 a	9.96 c	0.313 c	0.223 b
	Male group			
	Yield**	Protein**	Lysine**	Methionine**
HK	12.17 b	10.20 b	0.353 a	0.221 c
HM	11.57 b	10.95 a	0.338 b	0.239 a
HY	13.30 a	9.98 c	0.317 c	0.229 b
	Female by male group			
	Yield <sup>ns</sup>	Protein <sup>ns</sup>	lysine*	methionine**
HK × HK	10.92	10.02	0.361 ab	0.202 d
HM × HM	10.82	11.45	0.362 a	0.261 a
HY × HY	14.14	9.51	0.287 e	0.212 cd
HK × HM	11.39	10.81	0.331 c	0.218 cd
HK × HY	12.75	9.95	0.316 d	0.217 cd
HM × HK	12.15	10.79	0.367 a	0.240 b
HM × HY	13.01	10.48	0.348 b	0.257 a
HY × HK	13.44	9.78	0.332 c	0.220 c
HY × HM	12.46	10.6	0.321 cd	0.237 b

\* Significant at the 0.05 probability level, \*\* Significant at the 0.01 probability level, <sup>ns</sup> not significant

#### Mean grain yield, quality characteristics, lysine and methionine levels of the hybrids

Mean values of the investigated traits for all 56 diallel crosses and commercial checks across four environments are given in Table 6. Our study showed significant variation among genotypes ( $p < 0.01$ ) for grain yield and compositional traits. Grain yield ranged from 16.1 to 9.3 t ha<sup>-1</sup>. Normal × normal endosperm (HY × HY) hybrids including commercial checks had higher yield as expected. A1 × S1 (16.1 t ha<sup>-1</sup>) and M2 × S1 (15.5 t ha<sup>-1</sup>) were superior to commercial check means (14.7 t ha<sup>-1</sup>). The high grain yield of M2 × S1 (HM × HY) showed that high grain yield can be achieved in *dzr1* × normal endosperm crosses. On the other hand, in general lower grain yields were obtained from crosses containing high amino acid inbreds.

The mean protein values of the experimental crosses and checks were 10.4 % and 9.5 % respectively. The highest protein values were obtained from M2 × M1 (12.1 %), M1 × K1 (11.6 %) and M1 × A1 (11.6 %) showing that *dzr1* germplasm combined well in terms of protein concentration. Whole kernel lysine concentration varied from 0.26 g/100g dry matter (S2 × A1) to 0.40 g/100g dry matter (K1 × M1) in hybrids. The mean of the crosses (0.33 g/100 g) was higher than mean of checks (0.29 g/100g). The M2 × M3, M2 × A1, M2 × S1, M3 × K1, M3 × K2 hybrid combinations all had 0.38 g/100 g lysine showing that *dzr1* (M2 and M3) inbred lines performed well for lysine concentration. The lowest lysine values were generally obtained from non-*opaque-2* and non-*dzr1* germplasm demonstrating that at least one parent must contain *opaque-2* or *dzr1* to get a high lysine hybrid. Scott et al. (2009) compared normal genotypes with high lysine genotypes that gave 0.290 g / 100g and 0.330 g / 100 g lysine, respectively. The values in high-lysine-containing

genotypes ranged from 0.390 to 0.510 g/100g. In a study of Carena and Dong (2017), experiment averages were found to be 0.318 g/100 g and 0.303 g / 100 g for trial 1 and 2, respectively. Findings of our study were slightly lower than Scott et al. (2009). However, our results were consistent with Carena and Dong (2017), with relatively higher results. Different genotypes and different environments and methods were used in the studies and could explain the differences observed.

Methionine values of the hybrids ranged from 0.28 to 0.19 g/100 g and the mean of crosses (0.23 g/100g) was significantly higher than the mean of checks. M2 × S1 (0.28 g/100g), M1 × M2 (0.27 g/100g), M2 × M1 (0.27 g/100g) and M2 × S2 (0.27 g/100g) were the best hybrids for methionine. Kernel methionine results showed that *dzr1* containing lines were successful at producing high methionine hybrids. Darrigues et al. (2005) reported a methionine concentration of 0.210 g / 100 g, in the *dzr1* inbred B101. Huffman et al. (2016) found this value to be 0.179 g / 100 g in hybrids in diallel crosses in which some parents contained *dzr1*. Although the findings were similar, relatively higher results were obtained from our study. A microbial method was used to determine the amount of methionine in Darrigues et al (2005) and Huffman et al. (2016). In our study, a high resolution chromatographic (LC-MS / MS) method was used. Differences in the the methods used may be partially responsible for the observed differences among studies.

LQI values ranged from 3.9 to 2.6% and the best hybrid combinations were M3 × K1 (3.9 %), M2 × S1 (3.8 %) and K1 × M1 (3.7%) hybrids. MQI values varied from 1.8 % (K1 × M2) to 2.7% (M2 × S1). The overall experiment average was 2.3% and the most successful hybrids were M2 × S1 (2.7%) and M3 × S1 (2.6%) genotypes, respectively.

**Table 6.** Means of traits of the 56 hybrids obtained from an 8 × 8 full diallel mating design and commercial checks generated from two years and two sites

Hybrids	Yield (t ha <sup>-1</sup> )	Protein (%)	Lysine (g/100g)	Methionine (g/100g)	KQI (%)	MQI (%)	Lysine yield (kg ha <sup>-1</sup> )	Methionine yield (kg ha <sup>-1</sup> )
K1 × K2	11.5	10.0	0.37	0.19	3.6	1.9	42.2	22.0
K1 × M1	10.7	10.9	0.40	0.24	3.7	2.3	42.3	26.0
K1 × M2	12.4	10.6	0.28	0.19	2.6	1.8	34.4	23.0
K1 × M3	11.0	9.7	0.29	0.22	3.1	2.3	31.7	24.3
K1 × A1	12.8	9.4	0.29	0.22	3.1	2.4	37.1	27.9
K1 × S1	13.8	9.6	0.30	0.19	3.2	2.0	41.6	26.6
K1 × S2	12.0	10.3	0.35	0.23	3.5	2.3	41.7	27.3
K2 × K1	10.3	10.0	0.36	0.21	3.6	2.1	37.5	21.7
K2 × M1	10.0	11.1	0.37	0.23	3.2	2.0	36.4	22.8
K2 × M2	12.1	11.1	0.33	0.22	2.9	1.9	39.7	25.7
K2 × M3	12.0	11.5	0.33	0.21	2.9	1.8	39.7	25.5
K2 × A1	12.5	10.5	0.30	0.22	2.9	2.1	37.4	26.9
K2 × S1	13.6	10.1	0.32	0.21	3.2	2.1	43.8	28.8
K1 × S2	11.9	9.9	0.33	0.23	3.3	2.3	39.1	26.9
M1 × K1	12.4	11.6	0.37	0.24	3.1	2.1	45.6	30.3
M1 × K2	10.6	11.0	0.37	0.24	3.3	2.2	39.7	25.8
M1 × M2	9.3	11.4	0.35	0.27	3.0	2.4	32.1	25.0
M1 × M3	10.5	11.4	0.36	0.26	3.1	2.2	37.4	27.5
M1 × A1	10.8	11.6	0.34	0.24	2.9	2.1	36.5	26.2
M1 × S1	11.9	10.3	0.34	0.25	3.2	2.4	40.3	29.6
M1 × S2	11.0	10.8	0.30	0.26	2.8	2.4	33.8	28.6
M2 × K1	12.8	10.4	0.35	0.26	3.4	2.5	45.1	33.3
M2 × K2	11.4	11.5	0.35	0.25	3.0	2.1	40.1	29.1
M2 × M1	9.4	12.1	0.37	0.27	3.0	2.2	34.5	25.3
M2 × M3	12.6	11.5	0.38	0.29	3.3	2.5	48.3	36.3
M2 × A1	14.6	10.4	0.38	0.26	3.6	2.5	55.1	38.0
M2 × S1	15.5	10.0	0.38	0.28	3.8	2.7	58.6	42.5
M2 × S2	13.2	10.5	0.33	0.27	3.1	2.6	43.1	36.2
M3 × K1	12.0	9.8	0.38	0.23	3.9	2.4	45.7	27.4
M3 × K2	13.6	10.4	0.38	0.21	3.6	2.0	51.1	28.2
M3 × M1	11.3	11.3	0.34	0.23	3.0	2.1	38.2	26.3
M3 × M2	11.8	11.1	0.37	0.24	3.3	2.1	43.9	28.0
M3 × A1	14.5	11.0	0.37	0.25	3.5	2.4	54.3	36.8
M3 × S1	13.5	9.7	0.35	0.25	3.6	2.6	46.7	33.4
M3 × S2	12.0	10.0	0.35	0.25	3.5	2.5	41.4	29.4
A1 × K1	12.8	9.7	0.30	0.19	3.2	2.1	38.4	24.7
A1 × K2	13.9	10.1	0.36	0.22	3.6	2.2	50.6	30.8
A1 × M1	10.7	11.1	0.36	0.25	3.2	2.2	38.3	27.1
A1 × M2	13.0	10.9	0.36	0.24	3.3	2.2	47.3	32.0
A1 × M3	13.8	11.0	0.34	0.21	3.1	1.9	46.2	29.3
A1 × S1	16.1	9.6	0.30	0.20	3.3	2.2	47.6	32.4
A1 × S2	13.6	9.5	0.33	0.22	3.5	2.4	44.4	29.6
S1 × K1	14.5	9.4	0.32	0.22	3.5	2.4	47.0	32.3
S1 × K2	14.4	10.1	0.35	0.24	3.6	2.4	49.6	33.9
S1 × M1	11.5	10.3	0.30	0.21	3.0	2.1	34.4	24.5
S1 × M2	13.6	10.3	0.29	0.24	2.9	2.3	39.9	32.5
S1 × M3	13.6	10.2	0.29	0.24	2.9	2.4	39.0	33.1
S1 × A1	13.9	9.6	0.28	0.21	3.0	2.2	37.4	29.3
S1 × S2	13.6	9.5	0.29	0.22	3.1	2.4	38.9	30.6
S2 × K1	12.2	9.6	0.32	0.23	3.5	2.4	39.0	27.8
S2 × K2	12.9	9.9	0.34	0.22	3.4	2.2	43.6	28.1
S2 × M1	11.6	10.9	0.31	0.24	2.9	2.2	35.9	28.4
S2 × M2	11.8	10.1	0.32	0.25	3.2	2.5	38.0	29.6
S2 × M3	12.6	10.7	0.32	0.24	3.1	2.3	40.5	31.0
S2 × A1	13.6	9.4	0.26	0.21	2.8	2.3	34.8	28.7
S2 × S1	14.1	9.5	0.28	0.21	3.0	2.2	38.9	29.5
P31G98	14.1	9.5	0.29	0.22	3.2	2.4	41.6	30.9
DKC6589	15.3	9.6	0.28	0.21	3.0	2.2	43.5	31.7
Mean of experiment	12.6	10.4	0.33	0.23	3.2	2.3	41.6	29.1
Mean of checks	14.7	9.5	0.29	0.21	3.1	2.3	42.5	31.3
Mean of crosses	12.5	10.4	0.33	0.23	3.2	2.3	41.6	29.0
CV	14.1	5.2	3.93	5.66	3.8	5.8	13.99	14.08
LSD	1.43**	0.43**	0.0129**	0.013**	0.12**	0.13**	4.7**	3.3**

\* Significant at the 0.05 probability level, \*\* Significant at the 0.01 probability level, <sup>ns</sup> not significant

Germplasm carrying *opaque-2* and *dzt1* has potential to increase grain lysine and methionine concentration. However, this germplasm may contribute to lower grain

yield. While grain value is normally determined by the mass of grain produced per area of land, an alternative value calculation could be based on the amount of amino

acid produced per area of land. Therefore, we calculated hybrid lysine and methionine yield (kg ha<sup>-1</sup>). Although cross and experiment lysine yield mean (41.6 kg ha<sup>-1</sup>) were not significantly different than check means (42.5 kg ha<sup>-1</sup>), 8 hybrids were significantly higher than the checks in this trait. Lysine yields ranged from 58.6 to 31.7 kg ha<sup>-1</sup>. M2 × S1 hybrid (58.6 kg ha<sup>-1</sup>) had 27 % more lysine yield than commercial high yielding normal hybrids. Methionine yield ranged from 42.5 to 22 kg ha<sup>-1</sup>. The best methionine yield hybrid, M2 × S1 (42.5 kg ha<sup>-1</sup>), was higher than the check mean by 26 % and 7 hybrids were significantly higher than the checks. Interestingly, the best lysine yield hybrids were also the best methionine yield hybrids.

Combinations of major genes have been used extensively in the development of sweet corn (Boyer and Shannon, 1983) or in the development of maize with novel starches (Wang et al., 1993). This study contributed information on methionine and lysine genetics.

In conclusion, the results suggest that high grain yield can be achieved in normal × normal or *dzr1* × normal endosperm crosses. In general, lower grain yields were obtained from crosses containing high amino acid inbreds. High lysine and methionine yields showed that combinations of *opaque-2* and *dzr1* might be useful in breeding programs with the goal of producing high yielding and high protein quality hybrids.

#### ACKNOWLEDGMENTS

This research was supported by The Scientific and Technological Research Council of Turkey (TUBITAK) (Project No: 1150394).

#### LITERATURE CITED

- AOAC International. 2002. Crude Protein in Cereal Grains and Oilseeds, Official Methods of Analysis of AOAC International. AOAC International, Washington, DC.
- Betran, F.J., A. Bockholt, F. Fojt, L. Rooney, R. Waniska. 2003a. Registration of parental lines: Registration of T×802. *Crop Sci.* 43 (5): 1891–1892.
- Betran, F.J., A. Bockholt, F. Fojt, G. Odvody. 2003b. Registration of T×807 maize line. *Crop Sci.* 43 (5):1892–1893.
- Boyer, C.D. and J.C. Shannon. 1983. The use of endosperm genes for sweet corn improvement. In: *Plant Breeding Reviews*, ed, Janick J. Springer, Boston, MA
- Carena, M.J. and N. Dong. 2017. The ND early QPM program: developing the next generation of healthier maize (*Zea mays* L.) products. *Euphytica* 213 (7):150.
- Chan, P.T. and P. Matanjun. 2017. Chemical composition and physicochemical properties of tropical red seaweed, *Gracilaria changii*. *Food Chem.* 221: 302-310.
- Darrigues, A., C. Buffard, K.R. Lamkey, M.P. Scott. 2005. Variability and genetic effects for tryptophan and methionine in commercial maize germplasm. *Maydica.* 50 (2):147–156.
- Duraes, F.O.M., P.C. Magalhaes, A.C. Oliveira, M.X. Santos, E.E. Gomes, C.T. Guimares. 2002. Combining ability of tropical maize inbred lines under drought stress conditions, *Crop Breed Appl Biotechnol.* 2 (2):291-298.
- Erdal, S. 2019. Determination of drought tolerance levels of maize inbred lines. *Turkjans.* 6 (2): 178–189.
- Erdal, S., M. Pamukcu, A. Ozturk, K. Aydinsakir, S. Soylu. 2015. Combining abilities of grain yield and yield related traits in relation to drought tolerance in temperate maize breeding. *Turk J Field Crops.* 20 (2): 203-212.
- Faountoulakis, M. and H.W. Lahm. 1998. Hydrolysis and amino acid composition analysis of proteins. *J Chromatogr A.* 826 (2): 109-134.
- Fisher, R.A. 1935. *The Design of Experiments.* 7th ed. London: Oliver and Boyd; New York: Hafner.
- Griffing, B. 1956. Concept of general and specific combining ability in relation to diallel crossing systems. *Aust J Biol Sci.* 9 (4):463–493.
- Hallauer, A.R. and M.J. Carena. 2009. Maize breeding. In: *Handbook of Plant Breeding: Cereals*, ed, Carena M.J. Springer, New York, pp 3–98.
- Harms, R.H. and G.B. Russell. 1998. Layer performance when returned to a practical diet after receiving an amino acid-deficient diet. *J Appl Poultry Res.* 7 (2):175-179.
- Huffman, R.D., J.W. Edwards, L.M. Pollak, M.P. Scott. 2016. Interaction of genetic mechanisms regulating methionine concentration in maize grain. *Crop Sci.* 56 (5):1–11.
- Krivanek, A., H. Groote, N. Gunaratna, A. Diallo, D. Freisen. 2007. Breeding and disseminating quality protein maize for Africa. *Afr J Biotechnol.* 6 (4): 312-324
- Mbuya, K., K.K. Nkongolo, A. Kalonji-Mbuyi. 2011. Nutritional analysis of quality protein maize varieties selected for agronomic characteristics in a breeding program. *Int J Plant Breed Genet.* 5 (4): 317-327.
- Mertz, E.T., L.S. Bates, O.E. Nelson. 1964. Mutant that changes protein composition and increases lysine content of maize endosperm. *Science* 145 (3629):279–280.
- Nas, L., M. Lima, R. Vencovsky, P.B. Gallo. 2000. Combining ability of maize inbred lines evaluation in the three environment in Brazil. *Scientia Agricola* 57 (1): 129-134.
- Olsen, M.S., T.L. Krone, R.L. Phillips. 2003. BSSS53 as a donor source for increased whole-kernel methionine in maize: Selection and evaluation of high-methionine inbreds and hybrids. *Crop Sci.* 43 (5):1634–1642.
- Phillips, R.L., J. Suresh, M. Olsen, T. Krone. 2008. Registration of high-methionine versions of maize inbreds A 632, B73 and Mo17. *J Plant Reg.* 2(3):243-245.
- Phillips, R.L. and B.A. McClure. 1985. Elevated protein-bound methionine in seeds of a maize line resistant to lysine plus threonine. *Cereal Chem.* 62 (3):213–218.
- Ravindran, V. 2012. Poultry Feed Availability and Nutrition in Developing Countries. Food and Agriculture Organization of the United Nations, Poultry Development Review 1-4.
- Rodriguez, F., G. Alvarado, A. Pacheco, J. Crossa, J. Burgueño. 2015. AGD-R (Analysis of Genetic Designs with R for Windows) Version 2.0. <http://hdl.handle.net/11529/10202> International Maize and Wheat Improvement Center (Accessed September 15, 2017).
- Saki, A.A., H.R. Naseri, M.M. Tabatabaei, P. Zamani, M. Haghight. 2012. Estimates of methionine and sulfur amino acid requirements for laying hens using different models. *Braz J Poultry Sci.* 14 (3):159-232.
- Schutte, J.B. and J. De Jong. 1999. Ideal amino acid profile for poultry. In: *Feed Manufacturing in the Mediterranean Region. Recent Advances in Research and Technology*, ed, Brufao, J. and Tacon, A. Zaragoza: CIHEAM - Options Mediterraneennes. 259-263
- Scott, M.P., J.M. Peterson, A.R. Hallauer. 2009. Evaluation of combining ability of quality protein maize derived from U.S. public inbred lines. *Maydica* 54 (4): 449-456
- Scott, M.P., S. Bhatnagar and J. Betran. 2004. Tryptophan and methionine levels in quality protein maize breeding germplasm. *Maydica* 49 (2004):303–311.

- Scott, P.M., A. Darrigues, S.S. Timothy, K. Lamkey. 2008. Recurrent selection to alter grain methionine concentration and improve nutritional value of maize. *Crop Sci.* 48 (5):1705–1713.
- Steel, R.G.D. and J.H. Torrie. 1980. Principles and procedures of statistics. A biometrical approach, 2nd Edition, McGraw-Hill Book Company, New York.
- TTSM 2018. Maize. Technical instructions for agricultural trials in maize. [https://www.tarimorman.gov.tr/BUGEM/TTSM/Belgeler/Te%20scil/Teknik%20Talimatlar/S%C4%B1cak%20%C4%B0klim%20Tah%C4%B1llar%C4%B1/MISIR\\_TEKNIK\\_TALIMA%20TI.pdf](https://www.tarimorman.gov.tr/BUGEM/TTSM/Belgeler/Te%20scil/Teknik%20Talimatlar/S%C4%B1cak%20%C4%B0klim%20Tah%C4%B1llar%C4%B1/MISIR_TEKNIK_TALIMA%20TI.pdf) (Accessed February 15, 2020, in Turkish).
- Vasal, S.K., G. Srinivasan, S. Pandey, C.F. Gonzalez, J. Crossa, D.L. Beck. 1993. Heterosis and combining ability of CIMMYT's quality protein maize germplasm I: Lowland tropical. *Crop Sci.* 33 (1):46–51.
- Vivek, B.S., A.F. Krivanek, N. Palacios, S.T. Afriyie, A.O. Diallo. 2008. Breeding Quality Protein Maize (QPM): Protocols for Developing QPM Cultivars. Mexico, D.F: CIMMYT.
- Wang, Y.J., P. White, L. Pollak and J. Jane. 1993. Characterization of starch structures of 17 maize endosperm mutant genotypes with Oh43 inbred line background. *Cereal Chem.* 70 (2): 171-178.