

## EFFECT OF FOLIAR APPLICATION OF BIOSTIMULANTS ON FORAGE YIELD IN ALFALFA (*Medicago sativa* L.)

Diana MARINOVA<sup>1\*</sup> , Svetlana STOYANOVA<sup>1</sup> , Iliyana PETROVA<sup>2</sup> 

<sup>1</sup>Institute of Agriculture and Seed Science "Obraztsov chiflik", Agricultural Academy, Bulgaria

<sup>2</sup>Institute of Cryobiology and Food Technology, Agricultural Academy, Bulgaria

\*Corresponding author: diana27hm@abv.bg

Received: 21.10.2022

### ABSTRACT

The present study was aimed to determine the effect of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> experimental biostimulants application on yield related main traits (plant height and stems number per m<sup>2</sup>) and on forage productivity of Prista 4 alfalfa variety. The study was carried out in experimental field at Institute of Agriculture and Seed Science "Obraztsov chiflik" - Ruse, from 2019 to 2021. The biostimulants were foliarly applied two times in each regrowth. There was found positive effect of the biostimulants application on the growth and development of alfalfa. The additional introducing of VR<sub>10</sub> and VR<sub>1</sub> biostimulants contributed to higher alfalfa grass stands by 17.13 and 14.36%, respectively compared to control. Foliar spraying with V<sub>2</sub> biostimulant had more clear pronounced positive effect on the stem development. According to the three-year results the average stems number per m<sup>2</sup> at V<sub>2</sub> treatment significant exceeded the control by 4.78%. V<sub>2</sub> biostimulant distinguished with a stronger positive effect on the forage productivity among studied products. Its application resulted in significant increases on green mass and dry matter yields by 4.18 and 4.99%, respectively over the control. V<sub>2</sub> application gives promising results and reason to be recommended as a corrective component in technology of growing alfalfa for forage.

**Keywords:** Alfalfa, biostimulants, plant height, stems number, yield.

### INTRODUCTION

The satisfaction of the population plant and animal productions needs poses a serious challenge to agriculture (Santini et al., 2021). A high yield per unit area obtaining is based on the cultivation of varieties with high genetic potential using appropriate agronomic practices.

Inorganic products (synthetic fertilizers and pesticides) application is an effective method to increase yields, but is costly (30 to 50% of production costs, depending on the crop) and also leads to environmental problems (Kopittke et al., 2019; Zambrano-Mendoza et al. 2021).

Innovative environmentally friendly practices, considering aspects of soil and groundwater health and biodiversity conservation, are the reduction of the use of chemical products and application of technologies for organic production (Mehmed and Enchev, 2020).

The biobased products such as organic foliar fertilizers, growth regulators and biostimulants represent a sustainable, efficient technology or complement to their synthetic counterparts (i.e., agrochemicals) to improve nutrient use efficiency and secure yields stability of agriculture under optimal and stressful conditions

(Bozhanska et al., 2017; Mihova et al., 2017; Stoyanova et al., 2021).

The foliar fertilization has been the subject of many field studies and has become widely adopted as a standard practice for many crops in a lot of countries (Ozturk and Yildirim, 2013; Wani et al., 2017; Ivanova-Kovacheva et al., 2018; Ivanov and Nikolov, 2021).

The alfalfa (*Medicago sativa* L.), also known as "Queen of Forage" is the most important perennial forage legume widely grown throughout the world because its high yield, quality, nutritional value and adaptability (Bouton, 2012). In Europe alfalfa is considered to be the species with the greatest contribution to sustainability of the agriculture (Huyghe, 2003). The yield increasing and improving the quality of alfalfa is of great importance for protein problem deciding under now existing market relations. A number of studies are related to search opportunities for increasing alfalfa forage yield by applying different agrotechnical practices - optimal density; intensity of utilization; intercropping, balanced fertilization (Sevov et al., 2007; Ceylan et al., 2009; Petkovic et al., 2019; Basaran et al., 2020).

The fertilization of alfalfa has been traditionally focused on effects of direct application of phosphorus (P) and/or potassium (K) in various combinations (Berg et al., 2007, Macolino et al., 2013). There is a little data available, indicating that integrated alfalfa nutrition involving the combined use of inorganic and organic fertilizers increases yield more than they application alone (Quansah, 2000). Vasileva and Kostov (2015) have reported the alfalfa yield was more stable under organic fertilization conditions as compared with mineral ones.

The present study was aimed to determine the effect of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> experimental biostimulants application, an elaboration of the Institute of Cryobiology and Food Technologies (ICFT) - Sofia, on yield related main traits (plant height and stems number per unit area) and on forage productivity of alfalfa.

### MATERIAL AND METHODS

The study was carried out in experimental field at Institute of Agriculture and Seed Science (IASS) "Obraztsov chiflik" – Ruse, Agricultural Academy,

Bulgaria from 2019 to 2021. The soil type of experimental site was leached black soil, located on sandy clay. Active soil fertility was characterized by good potassium (33.17 mg 100 g<sup>-1</sup> soil), insufficient nitrogen (16.84 mg 100 g<sup>-1</sup> soil) and poor phosphorus (6.15 mg 100g<sup>-1</sup> soil) nutrient regime. The humus content was low and ranged from 2.03% to 2.17% (for the layer from 0 to 40 cm). The soil reaction was slightly acid (pH - from 5.84 to 5.94).

Data for monthly temperature sums and rainfall during the three growing seasons are presented in Table 1. Long-term norm (LTN) represents average value of collected data about monthly temperatures and precipitation in the period 1895-2005. There was a considerable variability in rainfall amounts and its distribution during years of study. The year of alfalfa sowing characterized by a high temperatures and sufficient soil moisture in March. The precipitation rate in April and May of first year of the experiment was above LTN by 25.9 mm and 32.2 mm, respectively. There were prolonged rainy periods in June and August. The monthly temperature sums were close to LTA, with higher temperatures in April.

**Table 1.** Monthly temperature sums and monthly precipitation from 2019 to 2021

Months	Monthly rainfall, mm				Σ T °C			
	2019	2020	2021	LTN*	2019	2020	2021	LTN*
Σ October- March	327.2	259.7	320.3	235,1	1191.2	758.9	837.6	
April	76.6	75.2	8.6	50.7	437.5	345.4	483.7	341.8
May	98.3	90.3	37.1	66.1	493.2	506.2	605.6	512.2
June	74.2	62.1	105.4	80.5	660.7	655.2	657.8	606.6
July	2.2	84.7	167.8	67.4	762.3	727.4	692.6	697.7
August	54.1	34.0	13.3	49.4	708.8	735.9	739.4	739.8
September	16.7	42.2	65.8	44.6	583.0	586.0	570.6	535.3
Σ April-September	322.1	388.5	398.0	358.7	3556.1	3507.6	3749.7	3426.2

Legend: \* LTN (Long term norm) - the period from 1896 to 2006 was used

The second year of the experiment (2020) was distinguished by the monthly temperature sums close to LTN, and with slightly higher temperatures in June and July. Regarding the precipitation, except for June and August, during other months, its amount was higher than the LTN.

There were conditions of prolonged drought with higher temperatures and precipitation amount significantly below the relevant LTNs during April and May in the third year. From June to July, monthly temperature sums were close to LTN and the precipitation rate was significantly above LTN.

The investigation was conducted with Prista 4 alfalfa variety, without irrigation. According fall dormancy (FD) scored scale Prista 4 variety is moderately dormant alfalfa (FD rating 5). The study was designed in a randomized block design in 4 replicates, at a individual plot size of 5 m<sup>2</sup>. Sowing was performed by machine at a inter-row spacing of 12.5 cm with the seeding rate of 2.5 kg da<sup>-1</sup>. The field trial included a control (without foliar fertilization) and 3 treatments with V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> experimental biostimulants application. The biostimulants

were applied by foliar spraying two times in each regrowth at 0.3 l da<sup>-1</sup> application rate per treatment for all biostimulants. The first application was carried out at a plant height of 10-15 cm and the second one in alfalfa bud stage. The products applied were elaborated on the basis of 10% w/v vermicompost extracts and enrichment with an auxin-type growth regulator at the Laboratory of Biologically Active Substances to the ICFT - Sofia.

**V<sub>2</sub>** – combined vermicompost extracts with 2 extragents used, concentration 20% v/v.

**VR<sub>10</sub>** – vermicompost extract enriched with a regulator at a concentration of 10% v/v.

**VR<sub>1</sub>** – vermicompost extract enriched with a regulator at a concentration of 1% v/v.

During the foliar fertilization, the plants on the control treatment were treated with distilled water.

To examine the effect of biostimulants were estimated the agronomical traits as follows: plants height and plant stand density, green mass yield, dry matter content in green mass and dry matter yield.

The plants height (cm) was recorded by measuring the length of the stems from the soil surface to the tip of the uppermost leaf of live shoots at 5 locations per each harvesting plot. Assessment of plant stand density, expressed by stem number (SN) per m<sup>2</sup> was determined by counting the stems in sampling plot (50 cm x 50 cm) at each harvesting plot for the variants. Alfalfa was harvested at early flowering stage. The green mass yield (GMY), t ha<sup>-1</sup> were measured in 2019, 2020, and 2021 by harvesting of each plot and weighing in the field. A total 10 cuttings were made (2019 - two cuts, 2020 and 2021 - four cuts) during the study period. For dry matter content determination (%) 200 g fresh vegetative mass was sampled before cutting for each variant. The samples were dried to constant weight in a drying chamber at 105°C, weighed and the percentage of dry matter was estimated. Dry matter yield (DMY), t ha<sup>-1</sup> was determined on the basis of green mass yield and the percentage of dry matter.

The experimental data were analysed by the One-way analysis of variance (ANOVA) and means were separated by Least Significant Differences (LSD) at 5% probability level. The statistical analysis was performed using the STATGRAPHICS PLUS software.

## RESULTS AND DISCUSSION

According to the results obtained at this study, there were different effects of biostimulants application on the measured traits ( $p < 0.05$ ) in regrowths and growing seasons.

Data demonstrate that over the year of alfalfa establishment biostimulants were the most effective in first cut impacting positively stems elongation (Table 2). The reported values in second regrowth were in range from 66.40 cm in VR<sub>10</sub> treatment to 62.80 cm in VR<sub>1</sub> at 60.12 cm in control. It was found in establishment year the real plant height at VR<sub>10</sub>, VR<sub>1</sub> and V<sub>2</sub> treatments were significantly increased by 39.44%, 32.09% and 20.40%, respectively in comparison to untreated stand. The tendency outlined for positive impact of the biostimulants on the degree of phenotypic expression of the trait was kept in the first production year. The highest real plant height (average for four cuts) was ascertained at VR<sub>10</sub> and VR<sub>1</sub> biostimulants application (60 and 59 cm, respectively) in 2020. The mean values differences both among biostimulant treatments and biostimulants and control were considerable ( $p < 0.05$ ). In contrast to the first and second experimental years, the results from the third year indicated no significant differences in real plant height. For three years the maximum mean canopy height of 64.13 cm was obtained in VR<sub>10</sub> biostimulant application, followed by VR<sub>1</sub> treatment (62.61 cm). Analysis of variance showed that over period of investigation mean real plant height was not significantly different among VR<sub>1</sub> and VR<sub>10</sub> treatments but was considerably higher ( $p < 0.05$ ) versus V<sub>2</sub> treatment (59.29 cm) and control (54.75 cm).

**Table 2.** Effect of foliar application of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> biostimulants on grass stand height

Year	Treatment	Real plant height by cuts, cm				Mean	Difference %
		I	II	III	IV		
2019	Control	30.00	60.12	-	-	45.06 d	
	V <sub>2</sub>	44.50	64.00	-	-	54.25 b	120.40
	VR <sub>10</sub>	59.25	66.40	-	-	62.83 a	139.44
	VR <sub>1</sub>	56.25	62.80	-	-	59.52 ab	132.09
	<b>LSD (0.05)</b>					5.56	
2020	Control	47.75	60.25	59.00	31.00	49.50 d	
	V <sub>2</sub>	52.50	68.50	68.00	31.00	55.00 c	111.11
	VR <sub>10</sub>	68.00	71.50	69.00	31.50	60.00 a	121.21
	VR <sub>1</sub>	66.50	71.00	68.25	30.25	59.00 b	119.19
	<b>LSD (0.05)</b>					2.25	
2021	Control	92.75	75.75	83.75	26.50	69.69	
	V <sub>2</sub>	93.00	73.75	81.25	26.50	68.63	98.48
	VR <sub>10</sub>	91.75	69.75	87.50	29.25	69.56	99.81
	VR <sub>1</sub>	95.00	74.75	83.25	24.25	69.31	99.45
	<b>LSD (0.05)</b>					4.64	
2019-2021	Control					54.75 c	
	V <sub>2</sub>					59.29 b	108.30
	VR <sub>10</sub>					64.13 a	117.13
	VR <sub>1</sub>					62.61 a	114.36
	<b>LSD (0.05)</b>					2.77	
Annual	Average						
2019		47.50 b	63.33 b			55.42 b	
2020		58.69 b	67.81 ab	66.06 b	30.94 a	41.58 b	
2021		93.13 a	73.50 a	83.94 a	26.63 b	69.30 a	

Means followed by same letter in the columns are not significantly different at  $P \leq 0.05$

The same trend was observed in other studies demonstrated that the application of biobased products positively influenced the canopy height (Marinova et al., 2019). Results of present study are in agreement with those of Getnet and Raja (2013) who reported that plant height increased significantly with addition of vermicompost. Sefaoglu et al. (2021) also reported that the application of organic and inorganic fertilizers alone and in combination created differences in the plant height in sunflower.

Data recorded on grass stand density, expressed by stems number (SN) per m<sup>2</sup>, were present in Table 3. The means of the trait revealed that the response to the biostimulants application varied in regrowths and years. The treating alfalfa plant stands with the biostimulants not exhibited sustainable higher productivity of stems per unit area compared to the control, over the three years of study, with exception of V<sub>2</sub> treatment. In the year of alfalfa establishment no significant effect of the biostimulants on the studied trait was identified; however, a tendency to increase the potential for stems development was observed after foliar spraying with biostimulants, especially at VR<sub>10</sub> application in the secant cut of first year. The extent of trait phenotypic expression in the second growing season was significantly (p<0.05)

affected by V<sub>2</sub> biostimulants application. In treatment with V<sub>2</sub> 476 stems per unit area were counted and 448 in a stand without fertilization. The treatment with VR<sub>10</sub> biostimulant in 2021 had the most beneficial effect on increasing the stems number by 8.38%, as compared to the control. The stimulating effect of the VR<sub>10</sub> product was observed in all regrowths. The reported excess for V<sub>2</sub> was 3.86%. In turn, means for three-year of research confirmed that the highest stems number per unit area was obtained in V<sub>2</sub> treatment, as compared to the other biostimulants. Also, the excess of 4.78% compared to the untreated stand was statistically significant (p<0.05). The results were in agreement with the previous studies that foliar fertilization with some organic products contributed to development of grass stands with better density (Marinova et al., 2021).

According to a number of studies bio-based products contribute to increase the rate of photosynthesis and enhance metabolism (Popov and Dzimotudis, 2007) and positively affect the yield structural components (Churkova, 2008). On the contrary, Hall et al. (2002) in a study on the effectiveness and profitability of foliar fertilizers and growth regulators applied in alfalfa found that none of the tested foliar applied products increased stem density.

**Table 3.** Effect of foliar application of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> biostimulants on grass stand density

Year	Treatment	Stem number per m <sup>2</sup> by cuts				Mean	Difference %
		I	II	III	IV		
2019	Control	164	195	-	-	179.50 ab	
	V <sub>2</sub>	148	200	-	-	185.00 ab	103.06
	VR <sub>10</sub>	148	238	-	-	193.00 a	107.52
	VR <sub>1</sub>	144	209	-	-	176.50 b	98.33
	<b>LSD (0.05)</b>					13.96	
2020	Control	504	489	480	319	448.00 b	
	V <sub>2</sub>	504	495	512	394	476.00 a	106.25
	VR <sub>10</sub>	390	393	412	401	399.00 c	89.06
	VR <sub>1</sub>	444	460	436	418	440.00 b	98.21
	<b>LSD (0.05)</b>					27.36	
2021	Control	434	412	391	290	381.75 b	
	V <sub>2</sub>	438	411	389	348	396.50 ab	103.86
	VR <sub>10</sub>	470	427	420	338	413.75 a	108.38
	VR <sub>1</sub>	429	390	399	329	386.75 b	101.31
	<b>LSD (0.05)</b>					24.01	
2019-2021	Control					336.42 b	
	V <sub>2</sub>					352.50 a	104.78
	VR <sub>10</sub>					335.25 b	99.65
	VR <sub>1</sub>					334.42 b	99.41
	<b>LSD (0.05)</b>					14.26	
Annual		Average					
2019		151.00 b	210.50 c			180.75 c	
2020		460.50 a	459.25 a	460.00 a	383.00 a	440.69 a	
2021		442.75 a	410.00 b	399.75 b	326.25 a	394.69 a	

Means followed by same letter in the columns are not significantly different at P ≤ 0.05

Data regarding green mass yield of the alfalfa stand under different treatments both in the year of crop establishment and two production years were given in

Table 4. The impact of foliarly applied products on alfalfa biomass was different over the three years of study. In the year of alfalfa establishment the use of biostimulants

resulted in increase of green mass yield. The treatment of alfalfa stands with the VR<sub>10</sub> biostimulant had the most beneficial effect on increasing the fresh biomass amount. The excess of 13.26%, over the control was significant. The application of V<sub>2</sub> and VR<sub>1</sub> biostimulants increased the green mass yield by 9.86% and 3.15%, respectively, but not significantly compared to control. In 2019 no significant differences ( $p < 0.05$ ) were found between treatments including V<sub>2</sub>, VR<sub>1</sub> and VR<sub>10</sub> biostimulants. Results indicated that the positive effect of foliar spraying with studied biostimulants on the main economically trait

was clearly expressed in first production year. It was found foliar spraying with VR<sub>10</sub> significant increased yield at all regrowths, as in the establishment year. The values reported shown that total annual green mass yield varied from 111.55 t ha<sup>-1</sup> at VR<sub>1</sub> application to 105.75 t ha<sup>-1</sup> for V<sub>2</sub>. It is evident the VR<sub>1</sub>, V<sub>2</sub> and VR<sub>10</sub> treatments resulted in significant ( $p < 0.05$ ) increases in yield by 20.59%, 18.11% and 14.32%, respectively compared to the control. Unlike the first and second alfalfa growing season, treatment with the biostimulants not contributed to higher green mass yield during the last production year.

**Table 4.** Effect of foliar application of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> biostimulants on green mass yield

Year	Treatment	Green mass yield by cuts, t ha <sup>-1</sup>				Total	Difference %
		I	II	III	IV		
2019	Control	5.75	6.32	-	-	12.07 b	
	V <sub>2</sub>	6.26	7.00	-	-	13.26 ab	109.86
	VR <sub>10</sub>	6.45	7.22	-	-	13.67 a	113.26
	VR <sub>1</sub>	5.75	6.70	-	-	12.45 ab	103.15
	<b>LSD (0.05)</b>					1.51	
2020	Control	29.95	25.05	29.40	8.10	92.50 b	
	V <sub>2</sub>	35.15	31.00	34.00	9.10	109.25 a	118.11
	VR <sub>10</sub>	30.90	31.00	34.40	9.45	105.75 a	114.32
	VR <sub>1</sub>	33.10	32.10	36.50	9.85	111.55 a	120.59
	<b>LSD (0.05)</b>					7.29	
2021	Control	35.80	33.95	28.90	2.50	101.15 a	
	V <sub>2</sub>	36.55	29.10	24.55	1.60	91.80 b	90.76
	VR <sub>10</sub>	35.10	30.40	26.70	2.30	94.50 b	93.43
	VR <sub>1</sub>	35.65	27.95	24.15	2.50	90.25 b	89.22
	<b>LSD (0.05)</b>					4.55	
2019-2021	Control					68.57 b	
	V <sub>2</sub>					71.44 a	104.18
	VR <sub>10</sub>					71.31 a	103.99
	VR <sub>1</sub>					71.42 a	104.15
	<b>LSD (0.05)</b>					2.29	
Annual		Average					
2019		6.05 c	6.81 b			6.43 c	
2020		32.28 b	29.79 a	33.58 a	9.13 a	26.19 a	
2021		35.78 a	30.35 a	26.08 b	2.23 b	23.61 b	

Means followed by same letter in the columns are not significantly different at  $P \leq 0.05$

In terms of forage productivity for three-year study was found that application of the experimental biostimulants contributed to higher mean annual green mass yield compared to control. The summarized data shown that positive effect of V<sub>2</sub>, VR<sub>1</sub> and VR<sub>10</sub> biostimulants on green mass yield was equalized (by increase of 4.18, 4.15 and 3.99 %, respectively).

The same trend has been demonstrated by other authors who noted that the bio-based products increase the amount of nitrogen in the soil and support the growth of plants, resulting in higher yields (Zaman et al., 2011; Caliskan et al., 2021). Javaad and Panwar (2013) also reported the use of vermicompost alone or in combination with other organic or mineral fertilizers is effective in increasing the productivity of various plants.

Dry matter content is a characteristic which play a key role in alfalfa hay yield. Data in the present study showed

the biostimulants foliar application had no effect on dry matter content in fresh biomass in year of alfalfa establishment (Table 5). In 2020 higher variability of the trait between treatments was observed. The values were in range from 26.25% to 23.75%, whit priority of the control. There was found V<sub>2</sub> and VR<sub>1</sub> application exhibited slight positive effect on dry matter accumulation in third growing season. The stimulating effect of the both products was the most clearly expressed in the first regrowth. V<sub>2</sub> and VR<sub>1</sub> application led to dry matter content increase by 5%, as compared to the control. The positive impact of the both biostimulants was kept in the second and third regrowths. Mean trait values for the three-year study shown V<sub>2</sub> and VR<sub>10</sub> biostimulants application contributed to slight increase in dry matter content by 0.42 and 0.25 percentage units, respectively.

**Table 5.** Effect of foliar application of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> biostimulants on dry matter content in alfalfa biomass

Year	Treatment	Dry matter content by cuts, %				Mean	Difference +/-
		I	II	III	IV		
2019	Control	30	28	-	-	29.00	
	V <sub>2</sub>	31	29	-	-	30.00	+ 1.00
	VR <sub>10</sub>	31	30	-	-	30.50	+ 1.50
	VR <sub>1</sub>	30	29	-	-	29.50	+ 0.50
2020	Control	25	23	30	27	26.25	
	V <sub>2</sub>	23	21	29	26	24.75	-1.50
	VR <sub>10</sub>	21	24	29	26	25.00	-1.25
	VR <sub>1</sub>	16	23	30	26	23.75	-2.50
2021	Control	24	26	26	31	26.75	
	V <sub>2</sub>	29	28	26	31	28.50	1.75
	VR <sub>10</sub>	25	28	25	31	27.25	0.50
	VR <sub>1</sub>	29	27	27	31	28.50	1.75
2019-2021	Control					27.33	
	V <sub>2</sub>					27.75	0.42
	VR <sub>10</sub>					27.58	0.25
	VR <sub>1</sub>					27.25	-0.08
Annual	Average						
2019		30.50 a	29.00 a			29.75 a	
2020		21.25 b	22.75 c	29.50 a	26.25 b	24.94 b	
2021		26.75 a	27.25 b	26.00 b	31.00 a	27.75 c	

Data regarding dry matter yield for the treatments including V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> biostimulants were one-way with those for green mass yield in the year of alfalfa establishment (Table 6). However, a more significant increases were found in DMY yield at the biostimulants application than in GMY compared to the control. VR<sub>10</sub> application distinguished with stronger positive impact on annual DMY, compared to the other tested biostimulants. It can be seen that in the first production year, alfalfa DMY changed significantly among biostimulant treatment in response to the significant variation in dry matter content, mainly in the first and second regrowths. It was observed that in terms of DMY the excesses above the control were significantly reduced in comparison the increases determined regarding GMY. This is most pronounced at VR<sub>10</sub> treatment (from increase of 20.59% in GMY to 7.97% in DMY). In 2021, an opposite trend was observed. It was found that despite the relatively lower green mass yields at biostimulants foliar application, the total annual dry matter yield was equal to that of untreated variant. Over the study period the effect of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> biostimulants was expressed in mean annual DMY of 18.79, 18.40 and 18.35 t ha<sup>-1</sup> at harvested yield of 17.90 t ha<sup>-1</sup> from the untreated stand.

Our findings confirmed statements of Kavut and Avcioğlu (2015) that dry matter content of forage crops is one of the dependable criteria of biomass production and high rate of dry matter content is mostly indicate a better adaptability and yield performance.

In 2019 (year of alfalfa establishment) sufficient soil moisture and high temperatures in March ensured good

seed germination and normal initial development of young alfalfa plants. The significant rainfall amounts in April and May and prolonged rainy periods in June and August led to delay of growth and development of the alfalfa grass stands. In the first year of study alfalfa achieved only two cuts. In 2020 and 2021 (first and second production year) alfalfa achieved four cuts. This explains the higher annual yields in the second and third year of study.

Data shown that the average plant height in the first and third cuts and average plant height in 2021 were statistically higher compared to those in 2019 and 2020. In second and third cuts and average for 2020 more number of stems were determined than those reported in 2019 and 2021 and differences were statistically significant.

The annual green mass yield in 2020 was statistically higher than those achieved in 2019 and 2021. Regarding dry matter yield statistically significant differences between the measured yields in 2020 and 2021 were not observed.

It can note the estimated agronomical traits was significantly influenced by year and weather conditions. Our results confirm the findings that the sustainable yields of crops are the result of the complex interaction between internal factors (variety genetic composition) (Tucak et al., 2014) and external factors (year, weather conditions, management) (Petkovic et al., 2019). According Kocira et al. (2020), effectiveness of biostimulants is determined not only the crop species, variety and environment but also by appropriate choice of preparations, their dose and method of application.

**Table 6.** Effect of foliar application of V<sub>2</sub>, VR<sub>10</sub> and VR<sub>1</sub> biostimulants on dry matter yield

Year	Treatment	Dry matter yield by cuts, t ha <sup>-1</sup>				Total	Difference, %
		I	II	III	IV		
2019	Control	1.73	1.77	-	-	3.50 c	
	V <sub>2</sub>	1.94	2.03	-	-	3.97 ab	113.43
	VR <sub>10</sub>	2.00	2.17	-	-	4.17 a	119.14
	VR <sub>1</sub>	1.73	1.94	-	-	3.67 bc	104.86
	<b>LSD (0.05)</b>					442.7	
2020	Control	7.48	5.76	8.82	2.18	24.25 b	
	V <sub>2</sub>	8.08	6.24	9.86	2.59	26.77 a	110.44
	VR <sub>10</sub>	6.49	7.44	9.97	2.46	26.36 a	108.75
	VR <sub>1</sub>	5.30	7.38	10.95	2.56	26.19 ab	107.97
	<b>LSD (0.05)</b>					2.08	
2021	Control	8.59	8.83	7.75	0.78	25.95 a	
	V <sub>2</sub>	10.60	8.15	6.38	0.50	25.63 ab	98.78
	VR <sub>10</sub>	8.78	8.51	6.68	0.71	24.68 b	95.12
	VR <sub>1</sub>	10.34	7.55	6.52	0.77	25.19 ab	97.07
	<b>LSD (0.05)</b>					1.25	
2016-2018	Control					17.90 b	
	V <sub>2</sub>					18.79 a	104.99
	VR <sub>10</sub>					18.40 ab	102.83
	VR <sub>1</sub>					18.35 ab	102.53
	<b>LSD (0.05)</b>					654.5	
Annual	Average						
2019	1.85 c		1.98 c		1.91 b		
2020	6.84 b		6.71 b		6.47 a		
2021	9.58 a		8.26 a		6.34 a		

Means followed by same letter in the columns are not significantly different at  $P \leq 0.05$

## CONCLUSION

The application of experimental biostimulants had a positive effect on the growth and development of alfalfa in the environmental conditions of IASS "Obraztsov chiflik". The additional introducing of VR<sub>10</sub> and VR<sub>1</sub> biostimulants contributed to higher alfalfa grass stands. The positive impact resulted in significant increase in plant height by 17.13 and 14.36%, respectively. Foliar spraying with V<sub>2</sub> biostimulant had more clear pronounced positive effect on the potential for stem formation and development of grass stands with better density. According to the three-year results the average stems number per m<sup>2</sup> at V<sub>2</sub> treatment significant exceeded the control by 4.78%. V<sub>2</sub> biostimulant distinguished with a stronger positive effect on the GMY and DMY among studied experimental products. The mean annual GMY and DMY at treatment with the V<sub>2</sub> were higher by 4.18 and 4.99%, respectively compared to the control and the increases were statistically significant. Biostimulants application could be an important agrotechnical measure for increasing plant height and stem number as yield related main traits. Foliar fertilization with V<sub>2</sub> biostimulant gives promising results and reason to be recommended as a corrective component in technology of growing alfalfa for green forage and hay.

## ACKNOWLEDGEMENTS

The research was carried out with the financial support of the Scientific Research Fund, Ministry of Education and Science, within the framework of the implementation

of the Project "Use of biostimulants in organic cultivation of agricultural crops - assessment of contributions to the bioeconomy", Contract No KP-06-H46 /6 of 27.11.2020

## LITERATURE CITED

- Basaran, U., E. Gulumsar, M. Copur Dogrusor and H. Mut. 2020. Improving yield and quality in spring-sown alfalfa with annual companion crops. *Turkish Journal of Field Crops* 25(2):138-146.
- Berg, W.K., S.M. Cunningham, S.M. Brouder, B.C. Joern, K.D. Johnson, J.B. Santini and J.J. Volenec. 2007. The long-term impact of phosphorus and potassium fertilization on alfalfa yield and yield components. *Crop Science* 47:2198-2209.
- Bouton, J.H. 2012. Breeding lucerne for persistence. *Crop and Pasture Science* 63:95-106.
- Bozhanska, T., B. Churkova and T. Mihova. 2017. Influence of growth regulators and bio-fertilizers on productivity of perennial legume forage grasses in conditions of the Central Balkan Mountains. *Journal of Balkan Ecology*. 20(2):135-143.
- Caliskan, S., M. S. Hashemi, M. Akkamis, R. I. Aytekin and M. Bedir. 2021. Effect of gibberellic acid on growth, tuber yield and quality in potatoes (*Solanum tuberosum* L.). *Turkish Journal of Field Crops* 26(2):139-146.
- Ceylan, S., H. Soya, B. Budak, H. Akdemir and B.C. Esetlili. 2009. Effect of zinc on yield and some related traits of alfalfa. *Turkish Journal of Field Crops* 14:136-143.
- Churkova, B. 2008. Influence of some biologically active substances over the productivity of seeds of bird's-foot-trefoil. *Plant Science*. 45(1):82-85.
- Getnet, M. and N. Raja. 2013. Impact of vermicompost on growth and development of cabbage, *Brassica oleracea* Linn. and their sucking pest, *Brevicoryne brassicae* Linn.

- (Homoptera: Aphididae). Res. J. Environ. Earth Sci. 5:104-112.
- Hall, M.H., R.C. Stout and W.S. Smiles. 2002. Effects of foliar fertilizers and growth regulators on alfalfa yield and quality. Online. Crop Management. doi:10.1094/CM-2002-0429-01-RS.
- Huyghe, C. 2003. Les fourrages et la production de protéines. In: Actes des Journées de l'Association Française pour la Production Fourragère (AFPF). Paris, France, 17-32.
- Ivanova-Kovacheva, G., G. Dyakova, R. Mincheva. 2018. Studying of changes in the catalase enzyme activity in Prista table vine cultivar treated with the micronutrient Lebosol-Kalium. Journal of Mountain Agriculture on the Balkans 21(6): 134-141.
- Ivanov, L. and P. Nikolov. 2021. Study of the influence of the combination of foliar fertilizers Aminoazole and Lebozol Zinc 700 SC, on maize hybrid Ruse 464. Journal of Mountain Agriculture on the Balkans 24(4):429-441.
- Javaad, S. and A. Panwar. 2013. Effect of biofertilizer, vermicompost and chemical fertilizer on different biochemical parameters of *Glycine max* and *Vigna mungo*. Recent Research in Science and Technology 5(1):40-44.
- Kavut, Y.T. and R. Avcioglu. 2015. Yield and quality performances of various alfalfa (*Medicago sativa* L.) cultivars in different soil textures in a mediterranean environment. Turkish Journal of Field Crops 20(1): 65-71.
- Kocira, A., J. Lamorska, R. Kornas, N. Nowosad, M. Tomaszewska, D. Leszczynska, K. Kozlowicz and S. Tabor. 2020. Changes in biochemistry and yield in response to biostimulants applied in bean (*Phaseolus vulgaris* L.). Agronomy 10:189.
- Kopittke, P.M., N.W. Menzies, P. Wang, B.A. McKenna and E. Lombi. 2019. Soil and the intensification of agriculture for global food security. Environment International, 132, 105078. <https://doi.org/10.1016/j.envint.2019.105078>
- Macolino, S., L.M. Lauriault, F. Rimi and U. Ziliotto. 2013. Phosphorus and potassium fertilizer effects on alfalfa and soil in a non-limited soil. Agronomy Journal. 105:1613-1618.
- Marinova, D., I. Ivanova and G. Ivanova-Kovacheva. 2019. Study on the impact of Aminobest organic product on morphological and economic traits in Prista 5 alfalfa (*Medicago sativa* L.) variety. Field Crop Studies 12(1):161-174.
- Marinova, D., I. Ivanova and S. Stojanova. 2021. Application of Total Care and Aminobest foliar fertilizers and their influence on the main quality traits in alfalfa (*Medicago sativa* L.). Journal of Mountain Agriculture on the Balkans. 24(4):109-129.
- Mehmed, A. and S. Enchev. 2020. Influence of organic foliar fertilizers on stevia development (*Stevia rebaudiana* B.). Journal of Mountain Agriculture on the Balkans, 23(4):230-242.
- Mihova, T., D. Georgiev, B. Brashlyanova and P. Ivanova. 2017. Influence of organic fertilization on the biochemical composition of fresh and dried fruits of japanese quince (*Chaenomeles* sp.). In: Proceedings of the VIII International Scientific Agriculture Symposium AGROSYM 2017, October 2017, Jahorina, Bosna and Herzegovina, pp. 1654-1659.
- Ozturk, G. and Z. Yildirim. 2013. Effect of bio-activators on the tuber yield and tuber size of potatoes. Turkish Journal of Field Crops 18(1): 82-86.
- Petkovic, K., M. Manojlovic, P. Lombnaes, R. Cabilovski, D. Krstic, Z. Loncaric and P. Lombnaes. 2019. Foliar application of selenium, zinc and copper in alfalfa (*Medicago sativa* L.) biofortification. Turkish Journal of Field Crops 24(1):81-90.
- Popov, N. and A. Dzimotudis. 2007. Study of the effect of "RENI" on the quality and productivity of garden peas. Journal of Central European Agriculture. 8(4):413-418.
- Quansah, C. 2000. Integrated soil management for sustainable agriculture and food security. A case study for four countries in W/Africa. FAO/RAF. 1:33-75.
- Santini, G., N. Biondi, L. Rodolfi and M.R. Tredici. 2021. Plant biostimulants from cyanobacteria: An emerging strategy to improve yields and sustainability in agriculture. Plants. 10:643. <https://doi.org/10.3390/plants10040643>
- Sefaoglu, F., H. Ozturk, E. Ozturk, M. Sezek, Z. Toktay and T. Polat. 2021. Effect of organic and inorganic fertilizers, or their combination on yield and quality components of oil seed sunflower in a semi-arid environment. Turkish Journal of Field Crops 26(1):88-95.
- Sevov, A., Hr. Yancheva, N. Popov and Hr. Kirchev. 2007. Investigation on the effect of preparation Reny on the productivity of alfalfa variety Mnogolistna 1. Field Crops Studies 4(2):337-341.
- Stoyanova, Sv., I. Petrova and D. Marinova. 2021. Influence of sowing dates and treatment with organic products on the development of winter oilseed rape (*Brassica napus*). Journal of Mountain Agriculture on the Balkans. 24(4):186-205.
- Tucak, M., S. Popovic, T. Cupic, G. Krizmanic, V. Spanic, B. Simic and V. Meglic. 2014. Agro-morphological and forage quality traits of selected alfalfa populations and their application in breeding. Turkish Journal of Field Crops 19(1):79-83.
- Vasileva, V. and O. Kostov. 2015. Effect of alfalfa grown for forage on soil fertility related to mineral and organic fertilization. Emirates J. Food and Agriculture. 27:687-686.
- Wani, M.Y., M.R. Mir, M.F. Baqual, K. Mehraj, T.A. Bhat and S. Rani. 2017. Role of foliar sprays in sericulture industry. Journal of Pharm. and Phytoch. 6(4):1803-1806.
- Zaman, A., A. Sarkar, S. Sarkar and W.P. Devi. 2011. Effect of organic and inorganic sources of nutrients on productivity, specific gravity and processing quality of potato (*Solanum tuberosum*). Indian Journal of Agricultural Sciences 81(12):1137-1142.
- Zambrano-Mendoza, J.L., C.A. Sangoquiza-Caiza, D.F. Campana-Cruz and C.F. Yanez-Guzman. 2021. Use of biofertilizers in agricultural production. In Technology in Agriculture; Available online: <https://www.intechopen.com/onlinefirst/76918>